

RESEARCH

Open Access



Some explicit identities on Changhee-Genocchi polynomials and numbers

Byung-Moon Kim¹, Joohee Jeong² and Seog-Hoon Rim^{2*} *Correspondence: shrim@knu.ac.kr²Department of Mathematics
Education, Kyungpook National
University, 80 Daehakro, Bukgu,
Daegu, 41566, S. Korea
Full list of author information is
available at the end of the article

Abstract

In this paper, we introduce a new family of functions, which is called the Changhee-Genocchi polynomials. We study some explicit identities on these polynomials, which are related to Genocchi polynomials and Changhee polynomials. Also, we represent Changhee-Genocchi polynomials by gamma and beta functions.

We also study some properties of higher-order Changhee-Genocchi polynomials related to Changhee polynomials and Daehee polynomials.

MSC: 05A10; 05A19; 11B68; 11S80**Keywords:** Euler polynomials; Changhee polynomials; Genocchi polynomials; Changhee-Genocchi numbers; beta and gamma functions

1 Introduction

The Genocchi polynomials are defined by the generating function (see [1, 2])

$$\frac{2t}{e^t + 1} e^{xt} = \sum_n G_n(x) \frac{t^n}{n!}. \quad (1)$$

When $x = 0$, $G_n = G_n(0)$ are called the Genocchi numbers. From (1) we see that

$$\begin{aligned} \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!} &= \left(\frac{2t}{e^t + 1} \right) e^{xt} = \left(\sum_{l=0}^{\infty} G_l \frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} x^m \frac{t^m}{m!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l} G_l x^{n-l} \right) \frac{t^n}{n!}. \end{aligned} \quad (2)$$

We consider Changhee-Genocchi polynomials defined by the generating function

$$\frac{2 \log(1+t)}{2+t} (1+t)^x = \sum_{n=0}^{\infty} CG_n(x) \frac{t^n}{n!}. \quad (3)$$

When $x = 0$, $CG_n = CG_n(0)$ are called the Changhee-Genocchi numbers.

The gamma and beta functions are defined by the following definite integrals:

$$\Gamma(\alpha) = \int_0^{\infty} e^{-t} t^{\alpha-1} dt, \quad \alpha > 0, \quad (4)$$

and

$$\begin{aligned} B(\alpha, \beta) &= \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} dt \\ &= \int_0^{\infty} \frac{t^{\alpha-1}}{(1+t)^{\alpha+\beta}} dt, \quad \alpha > 0, \beta > 0. \end{aligned} \quad (5)$$

From (4) and (5) we have (see [3])

$$\Gamma(\alpha + 1) = \alpha \Gamma(\alpha), \quad B(\alpha, \beta) = \frac{\Gamma(\alpha) \Gamma(\beta)}{\Gamma(\alpha + \beta)}. \quad (6)$$

We recall that the classical Stirling numbers of the first kind $S_1(n, k)$ and $S_2(n, k)$ are defined by the relations (see [4])

$$\begin{aligned} (x)_n &= \sum_{k=0}^n S_1(n, k) x^k \quad \text{and} \\ x^n &= \sum_{k=0}^n S_2(n, k) (x)_k, \end{aligned}$$

respectively. Here $(x)_n = x(x-1) \cdots (x-n+1)$ denotes the falling factorial polynomial of order n . We also have

$$\begin{aligned} \sum_{n=m}^{\infty} S_2(n, m) \frac{t^n}{n!} &= \frac{(e^t - 1)^m}{m!} \quad \text{and} \\ \sum_{n=m}^{\infty} S_1(n, m) \frac{t^n}{n!} &= \frac{(\log(1+t))^m}{m!}. \end{aligned} \quad (7)$$

In this paper, we introduce a new family of functions, which is called the Changhee-Genocchi polynomials.

We study some properties of these polynomials, which are related to Genocchi polynomials and Changhee polynomials. Also we represent Changhee-Genocchi polynomials by gamma and beta functions.

We also study higher-order Changhee-Genocchi polynomials related to Changhee polynomials and Daehee polynomials.

Most of the ideas in this paper come from Kim and Kim [5]. Specifically, equations (14), (21), and (22) are related to the papers [5–8].

2 Changhee-Genocchi polynomials

First, we relate our newly defined Changhee-Genocchi polynomials to Genocchi polynomials.

Replacing t by $e^t - 1$ in (3) and applying (7), we get

$$\begin{aligned} \frac{2t}{e^t + 1} e^{tx} &= \sum_{n=0}^{\infty} CG_n(x) \frac{1}{n!} (e^t - 1)^n \\ &= \sum_{n=0}^{\infty} CG_n(x) \frac{1}{n!} \sum_{k=n}^{\infty} S_2(k, n) \frac{t^k}{k!} \\ &= \sum_{k=0}^{\infty} \left(\sum_{n=0}^k CG_n(x) S_2(k, n) \right) \frac{t^k}{k!}. \end{aligned} \quad (8)$$

The left-hand side of (8) is the generating function of the Genocchi polynomials. Thus, by comparing the coefficients of (1) and (8) we have the following theorem.

Theorem 1 *For any nonnegative integer k , we have*

$$G_k(x) = \sum_{n=0}^k CG_n(x) S_2(k, n). \quad (9)$$

On the other hand, if we replace t by $\log(1+t)$ in (1) and apply (7), then we get

$$\begin{aligned} \frac{2 \log(1+t)}{2+t} (1+t)^x &= \sum_{n=0}^{\infty} G_n(x) \frac{1}{n!} (\log(1+t))^n \\ &= \sum_{n=0}^{\infty} G_n(x) \frac{1}{n!} \sum_{k=n}^{\infty} S_1(k, n) \frac{t^k}{k!} \\ &= \sum_{k=0}^{\infty} \left(\sum_{n=0}^k G_n(x) S_1(k, n) \right) \frac{t^k}{k!}, \end{aligned} \quad (10)$$

where $S_1(k, n)$ are the Stirling numbers of the first kind.

By comparing the coefficients of both sides of (10), we get the following theorem.

Theorem 2 *For any nonnegative integer k , we have*

$$CG_k(x) = \sum_{n=0}^k G_n(x) S_1(k, n). \quad (11)$$

Remark When $x = 0$ in (11), we can see that Changhee-Genocchi numbers are integers.

We can consider equation (11) as the inversion formula for (9). From (3) we can consider the following identity:

$$\begin{aligned} \sum_{n=0}^{\infty} CG_n(x) \frac{t^n}{n!} &= \frac{2 \log(1+t)}{2+t} (1+t)^x = \left(\sum_{l=0}^{\infty} CG_l \frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} (x)_m \frac{t^m}{m!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l} CG_l(x) \right) \frac{t^n}{n!}. \end{aligned} \quad (12)$$

Thus, by comparing the coefficients of both sides of (12) we have

$$\begin{aligned} CG_n(x) &= \sum_{l=0}^n \binom{n}{l} CG_l(x)_{n-l} = \sum_{l=0}^n \binom{n}{l} CG_{n-l}(x)_l \\ &= \sum_{l=0}^n \left(\sum_{m=0}^{n-l} \binom{n}{l} CG_l S_1(n-l, m) x^m \right). \end{aligned} \quad (13)$$

From (13) we can derive the following theorem.

Theorem 3 *For any nonnegative integer n , we have*

$$\int_0^1 CG_n(x) dx = \sum_{l=0}^n \sum_{m=0}^{n-l} \binom{n}{l} CG_l S_1(n-l, m) \frac{1}{m+1}. \quad (14)$$

In this paper, we define the λ -Changhee-Genocchi polynomials by a generating function as follows:

$$\frac{2 \log(1+t)}{(1+t)^\lambda + 1} (1+t)^{\lambda x} = \sum_{n=0}^{\infty} CG_{n,\lambda}(x) \frac{t^n}{n!}. \quad (15)$$

We recall that the λ -Changhee polynomials are defined in [9] by

$$\frac{2}{(1+t)^\lambda + 1} (1+t)^{\lambda x} = \sum_{n=0}^{\infty} Ch_{n,\lambda}(x) \frac{t^n}{n!}. \quad (16)$$

When $\lambda = 1$, Changhee-Genocchi polynomials are well-known Changhee polynomials, cf. [10–18]. In order to establish a reflexive symmetry on the Changhee-Genocchi polynomials, we consider the following:

$$\begin{aligned} \sum_{n=0}^{\infty} CG_n(1-x) \frac{t^n}{n!} &= \frac{2 \log(1+t)}{1+(1+t)} (1+t)^{1-x} = -\frac{2 \log(1+t)}{(1+t)^{-1}+1} (1+t)^{-x} \\ &= \sum_{n=0}^{\infty} CG_{n,-1}(x) \frac{t^n}{n!}. \end{aligned} \quad (17)$$

By comparing the coefficients of (17) we have the following theorem.

Theorem 4 *For $n \in \mathbb{N}$, we have*

$$CG_n(1-x) = CG_{n,-1}(x). \quad (18)$$

Thus, from (3) and (18) we have

$$\begin{aligned} \sum_{n=0}^{\infty} CG_n(-x + (1-y)) \frac{t^n}{n!} &= \frac{2 \log(1+t)}{2+t} (1+t)^{-x+(1-y)} \\ &= \frac{2 \log(1+t)}{2+t} (1+t)^{-x} (1+t)^{1-y} \end{aligned}$$

$$\begin{aligned}
&= \left(\sum_{m=0}^{\infty} CG_m(-x) \frac{t^m}{m!} \right) \left(\sum_{l=0}^{\infty} (1-y)_l (-x) \frac{t^l}{l!} \right) \\
&= \sum_{n=0}^{\infty} \left(\sum_{m=0}^n \binom{n}{m} CG_m(-x) (1-y)_{n-m} \right) \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \sum_{m=0}^n \sum_{k=0}^{n-m} \binom{n}{m} CG_m(-x) S_1(n-m, k) (1-y)^k. \quad (19)
\end{aligned}$$

By comparing the coefficients of (19) we have

$$CG_n(1-(x+y)) = \sum_{m=0}^n \sum_{k=0}^{n-m} \binom{n}{m} CG_m(-x) S_1(n-m, k) (1-y)^k. \quad (20)$$

On the other hand, by (5), (6), and (20) we have

$$\begin{aligned}
&\int_0^1 y^n CG_n(1-(x+y)) dy \\
&= \sum_{m=0}^n \sum_{k=0}^{n-m} \binom{n}{m} CG_m(-x) S_1(n-m, k) B(n+1, k+1) \\
&= \sum_{m=0}^n \sum_{k=0}^{n-m} \binom{n}{m} CG_m(-x) S_1(n-m, k) \frac{\Gamma(n+1)\Gamma(k+1)}{\Gamma(n+k+2)}. \quad (21)
\end{aligned}$$

Thus, by (18) and (21) we have the following identities, which relate the λ -Changhee-Genocchi polynomials, the Stirling numbers, and the beta and gamma polynomials:

$$\begin{aligned}
&\int_0^1 y^n CG_{n-1}(x+y) dy \\
&= - \sum_{l=0}^n \sum_{m=0}^{n-l} \binom{n}{l} S_1(n-l, m) CG_l \int_0^1 y^n (1-(x+y))^m dy \\
&= - \sum_{l=0}^n \sum_{m=0}^{n-l} \sum_{k=0}^m \binom{n}{l} \binom{m}{k} S_1(n-l, m) (-x)^{m-k} CG_l \int_0^1 y^n (1-y)^k dy \\
&= - \sum_{l=0}^n \sum_{m=0}^{n-l} \sum_{k=0}^m \binom{n}{l} \binom{m}{k} S_1(n-l, m) (-x)^{m-k} CG_l B(n+1, k+1) \\
&= - \sum_{l=0}^n \sum_{m=0}^{n-l} \sum_{k=0}^m \binom{n}{l} \binom{m}{k} S_1(n-l, m) (-x)^{m-k} CG_l \frac{\Gamma(n+1)\Gamma(k+1)}{\Gamma(n+k+2)}. \quad (22)
\end{aligned}$$

From (16) we consider

$$\begin{aligned}
\sum_{n=0}^{\infty} CG_{n,\lambda}(1-x) \frac{t^n}{n!} &= \frac{2 \log(1+t)}{(1+t)^\lambda + 1} (1+t)^{\lambda(1-x)} = \frac{2 \log(1+t)}{1 + (1+t)^{-\lambda}} (1+t)^{-\lambda x} \\
&= \sum_{n=0}^{\infty} CG_{n,-\lambda}(x) \frac{t^n}{n!}. \quad (23)
\end{aligned}$$

By comparing the coefficients of (23) we have the following theorem.

Theorem 5 For any nonnegative integer n , we have

$$CG_{n,\lambda}(1-x) = CG_{n,-\lambda}(x). \quad (24)$$

Remark If we take $\lambda = 1$ in Theorem 5, then we have the result in Theorem 4.

From the second line of (23) and from (16) we have

$$\begin{aligned} & \left(\sum_{l=1}^{\infty} \frac{(-1)^{l-1} t^l}{l} \right) \left(\sum_{m=0}^{\infty} Ch_{m,\lambda}(x) \frac{t^m}{m!} \right) \\ &= \sum_{n=1}^{\infty} \left(\sum_{l=1}^n \frac{(-1)^{l-1}}{l} \frac{Ch_{n-l,\lambda}(x)}{(n-l)!} n! \right) \frac{t^n}{n!}. \end{aligned} \quad (25)$$

By comparing the coefficients of (23) and (25) we have the following theorem.

Theorem 6 For any positive integer n , we have

$$CG_{n,\lambda}(x) = \sum_{l=1}^n \frac{(-1)^{l-1}}{l} Ch_{n-l,\lambda}(x) \frac{n!}{(n-l)!}.$$

For $r \in \mathbb{N}$, we define the Changhee-Genocchi polynomials $CG_n^{(r)}(x)$ of order r by the generating function

$$\left(\frac{2 \log(1+t)}{2+t} \right)^r (1+t)^x = \sum_{n=0}^{\infty} CG_n^{(r)}(x) \frac{t^n}{n!}. \quad (26)$$

From (26) we have the following relation between the Changhee-Genocchi polynomials of order r and the Changhee polynomials of order r :

$$\begin{aligned} & (\log(1+t))^r \left(\frac{2}{2+t} \right)^r (1+t)^x \\ &= \left(r! \sum_{l=r}^{\infty} S_2(l, r) \frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} Ch_m^{(r)}(x) \frac{t^m}{m!} \right) \\ &= \left(\sum_{l=0}^{\infty} S_2(l+r, r) \frac{r! t^{l+r}}{(l+r)!} \right) \left(\sum_{m=0}^{\infty} Ch_m^{(r)}(x) \frac{t^m}{m!} \right) \\ &= \left(\sum_{l=0}^{\infty} S_2(l+r, r) \binom{l+r}{r}^{-1} \frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} Ch_m^{(r)}(x) \frac{t^m}{m!} \right) t^r \\ &= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l} S_2(l+r, r) \binom{l+r}{r}^{-1} Ch_{n-l}^{(r)}(x) \right) \frac{t^{n+r}}{n!}. \end{aligned} \quad (27)$$

By comparing the coefficients of (26) and (27) we have the following theorem.

Theorem 7 For any nonnegative integer n , we have

$$CG_n^{(r)}(x) = \sum_{l=0}^n \binom{n}{l} \binom{l+r}{r}^{-1} S_2(l+r, r) Ch_{n-l}^{(r)}(x).$$

For $d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$, we have the following identity:

$$\sum_{a=0}^{d-1} (-1)^a (1+t)^a = \frac{1 + (1+t)^d}{2+t}. \quad (28)$$

So, for such $d \equiv 1 \pmod{2}$, from (28), (3), and (15) we see that

$$\begin{aligned} \sum_{n=0}^{\infty} CG_n(x) \frac{t^n}{n!} &= \frac{2 \log(1+t)}{2+t} (1+t)^x \\ &= \sum_{a=0}^{d-1} (-1)^a \frac{2 \log(1+t)}{(1+t)^d + 1} (1+t)^{d(\frac{a+x}{d})} \\ &= \sum_{a=0}^{d-1} (-1)^a \sum_{n=0}^{\infty} CG_{n,d} \left(\frac{a+x}{d} \right) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{a=0}^{d-1} (-1)^a CG_{n,d} \left(\frac{a+x}{d} \right) \right) \frac{t^n}{n!}. \end{aligned} \quad (29)$$

By comparing the coefficients in (29), for $d \equiv 1 \pmod{2}$, we have the following theorem.

Theorem 8 *For any nonnegative integer n and $d \equiv 1 \pmod{2}$, we have*

$$CG_n(x) = \sum_{a=0}^{d-1} (-1)^a CG_{n,d} \left(\frac{a+x}{d} \right). \quad (30)$$

We remark that, for $d \equiv 1 \pmod{2}$, from (9) and (30) we have the inversion of Theorem 8.

Theorem 9 *For any nonnegative integer n and $d \equiv 1 \pmod{2}$, we have*

$$\begin{aligned} G_k(x) &= \sum_{n=0}^k CG_n(x) S_2(k, n) \\ &= \sum_{n=0}^k \left(\sum_{a=0}^{d-1} (-1)^a CG_{n,d} \left(\frac{a+x}{d} \right) \right) S_2(k, n). \end{aligned}$$

From the generating function of the Changhee-Genocchi polynomials in (1), replacing t by $\lambda \log(1+t)$, we get

$$\begin{aligned} \frac{2\lambda \log(1+t)}{(1+t)^\lambda + 1} (1+t)^{\lambda x} &= \sum_{n=0}^{\infty} G_n(x) \frac{1}{n!} (\lambda \log(1+t))^n \\ &= \sum_{n=0}^{\infty} \lambda^n G_n(x) \left(\sum_{k=n}^{\infty} S_1(k, n) \frac{t^k}{k!} \right) \\ &= \sum_{k=0}^{\infty} \left(\sum_{n=0}^k \lambda^n G_n(x) S_1(k, n) \right) \frac{t^k}{k!}. \end{aligned} \quad (31)$$

Thus, the left-hand side of (31) can be represented by the λ -Changhee-Genocchi polynomials as follows:

$$\frac{2\lambda \log(1+t)}{(1+t)^\lambda + 1} (1+t)^{\lambda x} = \lambda \sum_{k=0}^{\infty} CG_{k,\lambda}(x) \frac{t^k}{k!}. \quad (32)$$

By comparing the coefficients of (31) and (32) we have the following theorem.

Theorem 10 *For any nonnegative integer k , we have*

$$CG_{k,\lambda}(x) = \sum_{n=0}^k \lambda^{n-1} G_n(x) S_1(k, n).$$

From the generating function of the Changhee-Genocchi numbers in (3) we want to see the recurrence relation for the Changhee-Genocchi numbers:

$$\begin{aligned} 2 \log(1+t) &= \sum_{n=0}^{\infty} CG_n \frac{t^n}{n!} (t+2) \\ &= \sum_{n=1}^{\infty} CG_n \frac{t^{n+1}}{n!} + \sum_{n=0}^{\infty} 2CG_n \frac{t^n}{n!} \\ &= \sum_{n=2}^{\infty} nCG_{n-1} \frac{t^n}{n!} + 2 \sum_{n=1}^{\infty} CG_n \frac{t^n}{n!} \\ &= 2CG_1 t + \sum_{n=2}^{\infty} (nCG_{n-1} + 2CG_n) \frac{t^n}{n!}. \end{aligned} \quad (33)$$

On the other hand, from the left-hand side of (33) we have

$$2 \log(1+t) = \sum_{n=1}^{\infty} (-1)^{n-1} 2(n-1)! \frac{t^n}{n!}. \quad (34)$$

By comparing the coefficients of (33) and (34) we have the following recurrence relation for the Changhee-Genocchi numbers.

Theorem 11 *We have*

$$\begin{aligned} CG_0 &= 0, \\ nCG_{n-1} + 2CG_n &= 2(n-1)!(-1)^{n-1} \quad \text{for } n \geq 1. \end{aligned}$$

From the higher-order Changhee-Genocchi polynomials

$$\left(\frac{2 \log(1+t)}{2+t} \right)^r (1+t)^x = \sum_{n=0}^{\infty} CG_n^{(r)}(x) \frac{t^n}{n!} \quad (35)$$

we can deduce

$$CG_0^{(r)}(x) = CG_1^{(r)}(x) = \cdots = CG_{r-1}^{(r)}(x) = 0. \quad (36)$$

Thus, from (36) we can rewrite (35) as follows:

$$\left(\frac{2\log(1+t)}{2+t}\right)^r (1+t)^x = \sum_{n=0}^{\infty} CG_{n+r}^{(r)}(x) \frac{t^{n+r}}{(n+r)!}. \quad (37)$$

We recall that the Dahee polynomials are defined by the generating function (see [9, 19])

$$\frac{\log(1+t)}{t} (1+t)^x = \sum_{n=0}^{\infty} D_n(x) \frac{t^n}{n!}.$$

When $x = 0$, $D_n = D_n(0)$ are called the Dahee numbers.

For $r \in \mathbb{N}$, the higher-order Dahee numbers are given by the generating function (see [9, 19, 20])

$$\left(\frac{\log(1+t)}{t}\right)^r = \sum_{n=0}^{\infty} D_n^{(r)}(x) \frac{t^n}{n!}.$$

From (28) we have

$$\begin{aligned} 2\log(1+t) \sum_{a=0}^{d-1} (-1)^a (1+t)^a &= \frac{2\log(1+t)}{2+t} + \frac{2\log(1+t)}{t+2} (1+t)^d \\ &= \frac{2\log(1+t)}{t} \left(\sum_{a=0}^{d-1} (-1)^a (1+t)^a \right) \\ &= \sum_{n=0}^{\infty} CG_n \frac{t^{n-1}}{n!} + \sum_{n=0}^{\infty} CG_n(d) \frac{t^{n-1}}{n!} \\ &= \sum_{n=0}^{\infty} \left(2 \sum_{a=0}^{d-1} (-1)^a D_n(a) \right) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\frac{CG_{n+1}}{n+1} + \frac{CG_{n+1}(d)}{n+1} \right) \frac{t^n}{n!}. \end{aligned} \quad (38)$$

Thus, from (38) we have the following theorem.

Theorem 12 For any nonnegative integer n and $d \equiv 1 \pmod{2}$, we have

$$2 \sum_{a=0}^{d-1} (-1)^a D_n(a) = \frac{CG_{n+1}}{n+1} + \frac{CG_{n+1,d}}{n+1}.$$

3 Changhee-Genocchi polynomials arising from differential equations

In this section, we give new identities on the Changhee-Genocchi numbers by using differential equations. We use the idea recently developed by Kwon et al. [21].

By equation (3) we can write the generating function for the Changhee-Genocchi numbers as follows:

$$F(t) = \frac{2\log(1+t)}{2+t} = \sum_{n=0}^{\infty} CG_n \frac{t^n}{n!}. \quad (39)$$

Let

$$G(t) = \log(1+t) \quad \text{and} \quad H(t) = \frac{2}{2+t}.$$

Then

$$\begin{aligned} G^{(N)}(t) &= \left(\frac{d}{dt}\right)^N G(t) = (-1)^{N-1}(N-1)!e^{-N \cdot G(t)}, \quad \text{and} \\ H^{(N)}(t) &= \left(\frac{d}{dt}\right)^N H(t) \\ &= \left(-\frac{1}{2}\right)^N N!e^{-(N+1) \cdot K(t)}, \quad \text{where } K(t) = \log(1+t/2). \end{aligned}$$

Thus,

$$\begin{aligned} F^{(N)}(t) &= \left(\frac{d}{dt}\right)^N F(t) = \sum_{k=0}^N \binom{N}{k} G^{(N-k)} H^{(k)} \\ &= \sum_{k=0}^N \binom{N}{k} (-1)^{N-k-1} (N-k-1)! e^{-(N-k)G(t)} \\ &\quad \times \left(-\frac{1}{2}\right)^k k! e^{-(k+1)K(t)} \\ &= \sum_{k=0}^N \binom{N}{k} (-1)^{N-1} \left(\frac{1}{2}\right)^k k! (N-k-1)! e^{-(N-k)G(t)} e^{-(k+1)K(t)}. \end{aligned} \quad (40)$$

On the other hand,

$$\begin{aligned} e^{-(N-k)G} e^{-(k+1)K} &= \left(\sum_{n=0}^{\infty} (-N+k)^n \frac{G^n}{n!}\right) \left(\sum_{l=0}^{\infty} (-(k+1))^l \frac{K^l}{l!}\right) \\ &= \left(\sum_{n=0}^{\infty} (-N+k)^n \sum_{m=n}^{\infty} S_1(m, n) \frac{t^m}{m!}\right) \\ &\quad \times \left(\sum_{l=0}^{\infty} (-(k+1))^l \sum_{j=l}^{\infty} \frac{1}{2^j} S_1(j, l) \frac{t^j}{j!}\right) \\ &= \sum_{m=0}^{\infty} \left(\sum_{n=0}^m (-N+k)^n S_1(m, n)\right) \frac{t^m}{m!} \\ &\quad \times \sum_{j=0}^{\infty} \left(\sum_{l=0}^j (-(k+1))^l S_1(j, l) \frac{1}{2^j}\right) \frac{t^j}{j!} \\ &= \sum_{s=0}^{\infty} \left(\sum_{m=0}^s \binom{s}{m} \sum_{n=0}^m (-N+k)^n S_1(m, n)\right) \\ &\quad \times \sum_{l=0}^{s-m} (-(k+1))^l S_1(s-m, l) \frac{1}{2^{s-m}} \frac{t^s}{s!}. \end{aligned} \quad (41)$$

From (39) we have

$$F^{(N)}(t) = \left(\frac{d}{dt}\right)^N F(t) = \sum_{m=0}^{\infty} CG_{N+m} \frac{t^m}{m!}. \quad (42)$$

By comparing the coefficients of (40), (41), and (42) we have new identities on the Changhee-Genocchi numbers as follows.

Theorem 13 *For any nonnegative integer s , we have*

$$CG_{s+N} = \sum_{m=0}^s \binom{s}{m} \left\{ \left(\sum_{n=0}^m (-N+k)^n S_1(m, n) \right) \left(\sum_{l=0}^{s-m} (-k-1)^l S_1(s-m, l) \frac{1}{2^{s-m}} \right) \right\} \\ \times \sum_{k=0}^N \binom{N}{k} (-1)^{N-1} \left(\frac{1}{2} \right)^k k! (N-k-1)!.$$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to this work. All authors read and approved the final manuscript.

Author details

¹Department of Mechanical System Engineering, Dongguk University, 123 Dongdae-ro, Gyeongju-si, Gyeongsangbuk-do, 38066, S. Korea. ²Department of Mathematics Education, Kyungpook National University, 80 Daehakro, Bukgu, Daegu, 41566, S. Korea.

Acknowledgements

The authors would like to express their sincere gratitude to the Editor, who gave us valuable comments to improve this paper.

Received: 25 June 2016 Accepted: 25 July 2016 Published online: 04 August 2016

References

- Kim, T: On the multiple q -Genocchi and Euler numbers. *Russ. J. Math. Phys.* **15**(4), 481-486 (2008)
- Srivastava, HM, Özarslan, MA, Kaanoğlu, C: Some generalized Lagrange-based Apostol-Bernoulli, Apostol-Euler and Apostol-Genocchi polynomials. *Russ. J. Math. Phys.* **20**(1), 110-120 (2013)
- Zill, DG, Cullen, MR: *Advanced Engineering Mathematics*. Jones & Bartlett, Sudbury (2006)
- Roman, S: *The Umbral Calculus*. Pure and Applied Mathematics, vol. 111, x+193 pp. Academic Press [Harcourt Brace Jovanovich, Publishers], New York (1984)
- Kim, DS, Kim, T: Some identities involving Genocchi polynomials and numbers. *Ars Comb.* **121**, 403-412 (2015)
- Kim, DS, Kim, T: A study on the integral of the product of several Bernoulli polynomials. *Rocky Mt. J. Math.* **44**(4), 1251-1263 (2014)
- Kim, T: Some properties on the integral of the product of several Euler polynomials. *Quaest. Math.* **38**, 553-562 (2015)
- Kim, T: A study on the q -Euler numbers and the fermionic q -integral of the product of several type q -Bernstein polynomials on \mathbb{Z}_p . *Adv. Stud. Contemp. Math. (Kyungshang)* **23**, 5-11 (2013)
- Kwon, HI, Kim, T, Seo, JJ: A note on degenerate Changhee numbers and polynomials. *Proc. Jangjeon Math. Soc.* **18**(3), 295-305 (2015)
- Ozden, H, Cangul, IN, Simsek, Y, Kurt, V: On the higher-order w - q -Genocchi numbers. *Adv. Stud. Contemp. Math.* **19**(1), 39-57 (2009)
- Jang, L-C, Ryoo, CS, Seo, JJ, Kwon, HI: Some properties of the twisted Changhee polynomials and their zeros. *Appl. Math. Comput.* **274**, 169-177 (2016)
- Kim, T: p -Adic q -integrals associated with the Changhee-Barnes' q -Bernoulli polynomials. *Integral Transforms Spec. Funct.* **15**(5), 415-420 (2004)
- Kim, T, Dolgy, DV, Kim, DS, Seo, JJ: Differential equations for Changhee polynomials and their applications. *J. Nonlinear Sci. Appl.* **9**, 2857-2864 (2016)
- Kim, DS, Kim, T: A note on Changhee polynomials and numbers. *Adv. Stud. Theor. Phys.* **7**(20), 993-1003 (2013)
- Kim, T, Kim, DS: A note on nonlinear Changhee differential equations. *Russ. J. Math. Phys.* **23**(1), 88-92 (2016)
- Kim, DS, Kim, T, Seo, JJ, Lee, SH: Higher-order Changhee numbers and polynomials. *Adv. Stud. Theor. Phys.* **8**(8), 365-373 (2014)
- Kim, T, Rim, S-H: New Changhee q -Euler numbers and polynomials associated with p -adic q -integrals. *Comput. Math. Appl.* **54**(4), 484-489 (2007)
- Rim, S-H, Pak, HK, Jeong, J, Kang, DJ: Changhee-Genocchi numbers and their applications. Submitted for publication

19. El-Desouky, BS, Mustafa, A: New results on higher-order Daehee and Bernoulli numbers and polynomials. *Adv. Differ. Equ.* **2016**, 32 (2016)
20. Wang, NL, Li, H: Some identities on the higher-order Daehee and Changhee numbers. *Pure Appl. Math.* **4**(5-1), 33-37 (2015)
21. Kwon, HI, Kim, T, Seo, JJ: A note on Daehee numbers arising from differential equations. *Glob. J. Pure Appl. Math.* **12**(3), 2349-2354 (2016)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
